

---

# Optimization of treatment methods for chronic diseases of the nose and paranasal sinuses combined with chronic dacryocystitis

*N.J. Khushvakova*

*F.A. Nurmukhammedov*

*U.F. Davronov*

Samarkand State Medical Institute Samarkand, Uzbekistan

Samarkand State Medical Institute Samarkand, Uzbekistan

Samarkand State Medical Institute Samarkand, Uzbekistan

The pathogenesis of the overgrowth of dacryorrhinostoma is based on many factors from the development of granulations and scars in the area of the bone opening to the development of the commissural opening, mucositis in the nasal cavity to purulent persistent forms of rhinosinusitis in the area nose-lacrimal duct. To improve the reparative processes after the operation of endoscopic dacriocystomy with the installation of silicone stent-drainage (from the intubation tube), a zinc hyalrate solution was used.

---

## References

1. Garashchenko M. V., Garashchenko T. I., Ilyenko L. I. "Elimination therapy in preventive programs for seasonal prophylaxis of influenza and SARS" Rus. honey. journal 2009.No.P. 52-56.
2. "Irrigation method for the treatment of allergic rhinitis" G. D. Tarasova and others // Ros.otorinolar. 2008. No. 2. P.105-108.
3. Kyulev A. I. "Irrigation therapy of the upper respiratory tract." M.: Medicine, 1987.125p.
4. Mokronosova M. A. "Irrigation therapy of the nasal cavity from the perspective of evidence-based medicine" Vestn. otorhinolar. 2009. No1.P. 51-53.
5. "Otorhinolaryngology: nat. Leadership »Ed. V.T. Palchun. M.: GEOTAR-Media, 2008. p. 173-187
6. Khushvakova N.Zh., Nurmukhammedov F.O. "Features of the treatment of patients with chronic dacryocystitis using hyaluronic acid preparations for endonasal dacryocystostomy" "Austria-science" journal. 2018.No 15. p. 8-10.