

Morphological changes of muscles of the crus and prevention of postoperative complications at patients with the diabetes mellitus at critical ischemia of the lower extremity

Bakhodir Barnoyevich Safoyev Bukhara State medical institute
Azizzhon Yakhyoevich Bukhara State medical institute
Rakhimov
Razhabboy Isroilovich Isroilov Tashkent medical Academy

The case history of 57 patients with critical lower limb ischemia in diabetes mellitus in the clinical database of the Bukhara State Medical Institute was studied. All examined patients were taken during surgery 2 grams of muscle tissue from m. soleus and m. gastrocnemius from one anatomical zone from the side of the amputated limb. Biopsy material was studied to determine the microflora and to predict the course of the wound process, and therefore the qualitative and quantitative content of the microbial bodies of the material was determined by bacteriological examination. The morphology of the soleus and biceps muscles was also studied. The results of the study showed a high growth of microbial bodies from the soleus muscle, the reason for this is necrosis of the muscle tissue, which was confirmed by the morphological study.

References

1. A.R coders. Prevention of pyoinflammatory complications of an amputating stump at patients with a diabetes mellitus. Дясс. candidate of medical sciences. Tajik state medical university of Abu Ali Ibni Sino. 2013.
2. Zinich E. L./Vysevayemost frequency, resistance and sensitivity of microflora of primary supplicative focuses to antibacterial drugs at patients with complicated сдс depending on its form. / Suchasni medichni technologist i i. No. 3-4, 2011. Str 128-131.
3. Mitish, V. A. Gnoyno - necrotic defeats of a neuroischemic form of a syndrome of diabetic foot. New opportunities of complex surgical treatment / V. A. Mitish, I. A. Eroshkin, A. V. Eroshenko//Endocrine surgery. - 2008. - No. 1. - Page 24-29
4. B. B. Safoyev, A. Ya. Rakhimov, M. S. Sharopova. / Microbiological assessment of tissue of muscles of a shin at amputation at patients of critical ischemia of the lower extremity //Tibbiyotda of a yanga of kuna. (NDM) of.2018-№2(22). Str 46-50.
5. A.V's pro-tires. The comparative characteristic of a wound process at patients with it is purulent - necrotic forms of a syndrome of diabetic foot//the International magazine of applied and basic researches. - 2010. - No. 12. - Page 52-54.
6. Aboyans V, Criqui MH, Abraham P, Allison MA, Creager MA,
7. Diehm C, et al. Measurement and interpretation of the ankle-brachial index: scientific statement from the American Heart Association. Circulation 2012; 126:2890-909.
8. B.B.Safoyev., A.Ya.Rakhimov...//Situation of the problems of diagnosis and treatment of the syndrome of diabetic foot in modern surgery. - New Day in Medicine. (NDM) of.2018-№1 (21). P. 48-55.
9. Rosenthal V.D. et al. International Nosocomial Infection Control Consortium report, data summary for 2002 - 2007, issued January 2008. Am J Infect Control Nov; 36(9): 627 — 637.

10. Pratik K. Dalal, MD, Anand Prasad, MD, FSCAI, RPVI. Contemporary Outcomes of Endovascular Intervention for Critical Limb Ischemia Department of Cardiovascular Diseases, University of Texas Health Science Center, 7703 Floyd Curl Drive, MC 7872, San Antonio, TX 78229, USA, <http://dx.doi.org/10.1016/j.iccl.2016.12.008> 2211-7458/17/zh 2017 Elsevier Inc.
11. Hirsch AT, Allison MA, Gomes AS, Corriere MA, Duval S, Ershow AG, et al. A call to action: women and peripheral artery disease: scientific statement from the American Heart Association. *Circulation* ; 125:1449-72.
12. ; 125:1449-72.
13. Nathan A. Johnson, Alex L. Barwick, Angela Searle, Martin J. Spink, Stephen M. Twigg, Vivienne H. Chuter. Self-reported physical activity in community-dwelling adults with diabetes and its association with diabetes complications. *Journal of Diabetes and Its Complications* 33 (2019) 33-38.